

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAEGS1646

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	MAY 01	New CAS web site launched
NEWS	3	MAY 08	CA/CAPLUS Indian patent publication number format defined
NEWS	4	MAY 14	RDISCLOSURE on STN Easy enhanced with new search and display fields
NEWS	5	MAY 21	BIOSIS reloaded and enhanced with archival data
NEWS	6	MAY 21	TOXCENTER enhanced with BIOSIS reload
NEWS	7	MAY 21	CA/CAPLUS enhanced with additional kind codes for German patents
NEWS	8	MAY 22	CA/CAPLUS enhanced with IPC reclassification in Japanese patents
NEWS	9	JUN 27	CA/CAPLUS enhanced with pre-1967 CAS Registry Numbers
NEWS	10	JUN 29	STN Viewer now available
NEWS	11	JUN 29	STN Express, Version 8.2, now available
NEWS	12	JUL 02	LEMBASE coverage updated
NEWS	13	JUL 02	LMEDLINE coverage updated
NEWS	14	JUL 02	SCISEARCH enhanced with complete author names
NEWS	15	JUL 02	CHEMCATS accession numbers revised
NEWS	16	JUL 02	CA/CAPLUS enhanced with utility model patents from China
NEWS	17	JUL 16	CAPLUS enhanced with French and German abstracts
NEWS	18	JUL 18	CA/CAPLUS patent coverage enhanced
NEWS	19	JUL 26	USPATFULL/USPAT2 enhanced with IPC reclassification
NEWS	20	JUL 30	USGENE now available on STN
NEWS	21	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS	22	AUG 06	BEILSTEIN updated with new compounds
NEWS	23	AUG 06	FSTA enhanced with new thesaurus edition
NEWS	24	AUG 13	CA/CAPLUS enhanced with additional kind codes for granted patents
NEWS	25	AUG 20	CA/CAPLUS enhanced with CAS indexing in pre-1907 records
NEWS EXPRESS	29	JUNE 2007: CURRENT WINDOWS VERSION IS V8.2, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 05 JULY 2007.	
NEWS HOURS	STN Operating Hours Plus Help Desk Availability		
NEWS LOGIN	Welcome Banner and News Items		
NEWS IPC8	For general information regarding STN implementation of IPC 8		

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 13:00:38 ON 25 AUG 2007

=> File .gerry2MBCE  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 13:01:09 ON 25 AUG 2007

FILE 'BIOSIS' ENTERED AT 13:01:09 ON 25 AUG 2007

Copyright (c) 2007 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 13:01:09 ON 25 AUG 2007  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 13:01:09 ON 25 AUG 2007  
Copyright (c) 2007 Elsevier B.V. All rights reserved.

=> S MIP-4 (L)CCRL2  
L1 1 MIP-4 (L) CCRL2

=> D ibib Abs l1

L1 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:547792 CAPLUS  
DOCUMENT NUMBER: 143:76842  
TITLE: Macrophage inflammatory protein-4 (MIP-4) as an endogenous ligand for CCRL2 and sequences of human MIP-4 and CCRL2

INVENTOR(S): Tinsley, Jonathon Mark  
PATENT ASSIGNEE(S): Oxagen Limited, UK  
SOURCE: PCT Int. Appl., 69 pp.  
CODEN: PIXXD2

DOCUMENT TYPE: Patent  
LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005057220	A2	20050623	WO 2004-GB5057	20041202
WO 2005057220	A3	20060202		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1692171	A2	20060823	EP 2004-801256	20041202
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK,			

BA, HR, IS, YU  
 JP 2007520210 T 20070726 JP 2006-542005 20041202  
 US 2007036781 A1 20070215 US 2006-579386 20060515  
 PRIORITY APPLN. INFO.: GB 2003-28275 A 20031205  
 GB 2004-3014 A 20040211  
 GB 2004-18568 A 20040819  
 WO 2004-GB5057 W 20041202

AB Macrophage inflammatory protein-4 (MIP-4; also known as DC-CK1, CCL18 and PARC) is identified as an endogenous ligand for CCRL2 (chemokine (C-C motif) receptor-like 2). The protein and cDNA sequences of human MIP-4 and CCRL2 are disclosed. Anti-CCRL2 antibody was blocking MIP-4 and synovial fluid induced monocyte chemotaxis. Anti-MIP-4 antibody was also blocking RA synovial fluid induced monocyte chemotaxis. This data demonstrates that MIP-4 is a major mediator of monocyte induced chemotaxis found in RA synovial fluid. CCRL2 modulators, such as antibodies against CCRL2 or MIP-4, is useful in treating an inflammatory disease, a disease associated with enhanced macrophage activity or an infection.

=> S Antibody(S)MIP-4 AND pd<=20041202

2 FILES SEARCHED...

L2 1 ANTIBODY(S) MIP-4 AND PD<=20041202

=> D ibib abs L2

L2 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1995:820779 CAPLUS

DOCUMENT NUMBER: 123:220290

TITLE: Cloning and therapeutic applications of human macrophage inflammatory proteins MIP-3, MIP-4, and MIP-1γ, or their antibodies or antagonists

INVENTOR(S): Li, Haodong; Rosen, Craig A.; Ruben, Steve; Adams, Mark D.

PATENT ASSIGNEE(S): Human Genome Sciences, Inc., USA

SOURCE: PCT Int. Appl., 63 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 10

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9517092	A1	19950629	WO 1994-US7256	19940628 <--
W: AU, CA, CN, JP, KR, NZ				
RM: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5556767	A	19960917	US 1993-173209	19931222 <--
US 5504003	A	19960402	US 1994-208339	19940308 <--
ZA 9403442	A	19951120	ZA 1994-3442	19940518 <--
CA 2179606	A1	19950629	CA 1994-2179606	19940628 <--
AU 9475497	A	19950710	AU 1994-75497	19940628 <--
AU 684539	B2	19971218		
EP 735818	A1	19961009	EP 1994-925671	19940628 <--
EP 735818	B1	20040331		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CN 1143894	A	19970226	CN 1994-194902	19940628 <--
JP 09506774	T	19970708	JP 1995-517397	19940628 <--
JP 3677288	B2	20050727		

JP 2002053490	A	20020219	JP 2001-196723	19940628 <--
AT 262914	T	20040415	AT 1994-925671	19940628 <--
PT 735818	T	20040730	PT 1994-925671	19940628 <--
ES 2214484	T3	20040916	ES 1994-925671	19940628 <--
CN 1321745	A	20011114	CN 2001-116577	20010416 <--
AU 777297	B2	20041007	AU 2002-15445	20020206 <--
US 2003147846	A1	20030807	US 2002-165233	20020610 <--
PRIORITY APPLN. INFO.:			US 1993-173209	A 19931222
			US 1994-208339	A 19940308
			JP 1995-517397	A3 19940628
			WO 1994-US7256	W 19940628
			US 1995-446881	B1 19950505
			US 1995-468775	B2 19950606
			AU 1997-46576	A3 19970930
			US 1999-334923	A3 19990617
			US 1999-334951	A3 19990617
			US 1999-334954	A3 19990617

AB There are disclosed human macrophage inflammatory protein-3, human macrophage inflammatory protein-4, and human macrophage inflammatory protein-1γ polypeptides and DNA (or RNA) encoding such polypeptides. There is also provided a procedure for producing such polypeptides by recombinant techniques and for producing antibodies against such polypeptides. In the invention there is also provided antagonist/inhibitors against such polypeptides which inhibit the functioning of such polypeptides. Another aspect of the invention provides a combination of the polypeptides of the present invention and a suitable pharmaceutical carrier for providing a therapeutically effective amount of the polypeptides for the treatment of various associated diseases.

=> S Antibody(S)CCRL-2 AND pd<=20041202

2 FILES SEARCHED...

L3 0 ANTIBODY(S) CCRL-2 AND PD<=20041202

=> Log Off h

SESSION WILL BE HELD FOR 120 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 13:06:52 ON 25 AUG 2007

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAEGS1646

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1		Web Page for STN Seminar Schedule - N. America
NEWS 2	AUG 06	CAS REGISTRY enhanced with new experimental property tags
NEWS 3	AUG 06	FSTA enhanced with new thesaurus edition
NEWS 4	AUG 13	CA/CaPlus enhanced with additional kind codes for granted patents
NEWS 5	AUG 20	CA/CaPlus enhanced with CAS indexing in pre-1907 records
NEWS 6	AUG 27	Full-text patent databases enhanced with predefined patent family display formats from INPADOCDB
NEWS 7	AUG 27	USPATOLD now available on STN

NEWS 8 AUG 28 CAS REGISTRY enhanced with additional experimental  
 spectral property data  
 NEWS 9 SEP 07 STN AnaVist, Version 2.0, now available with Derwent  
 World Patents Index  
 NEWS 10 SEP 13 FORIS renamed to SOFIS  
 NEWS 11 SEP 13 INPADOCDB enhanced with monthly SDI frequency  
 NEWS 12 SEP 17 CA/Caplus enhanced with printed CA page images from  
 1967-1998  
 NEWS 13 SEP 17 Caplus coverage extended to include traditional medicine  
 patents  
 NEWS 14 SEP 24 EMBASE, EMBAL, and LEMBASE reloaded with enhancements  
 NEWS 15 OCT 02 CA/Caplus enhanced with pre-1907 records from Chemisches  
 Zentralblatt  
 NEWS 16 OCT 19 BEILSTEIN updated with new compounds  
 NEWS 17 NOV 15 Derwent Indian patent publication number format enhanced  
 NEWS 18 NOV 19 WPIX enhanced with XML display format  
 NEWS 19 NOV 30 ICSD reloaded with enhancements  
 NEWS 20 DEC 04 LINPADOCDB now available on STN  
 NEWS 21 DEC 14 BEILSTEIN pricing structure to change  
 NEWS 22 DEC 17 USPATOLD added to additional database clusters  
 NEWS 23 DEC 17 IMSDRUGCONF removed from database clusters and STN  
 NEWS 24 DEC 17 DGENE now includes more than 10 million sequences  
 NEWS 25 DEC 17 TOXCENTER enhanced with 2008 MeSH vocabulary in  
 MEDLINE segment  
 NEWS 26 DEC 17 MEDLINE and LEMEDLINE updated with 2008 MeSH vocabulary  
 NEWS 27 DEC 17 CA/Caplus enhanced with new custom IPC display formats  
 NEWS 28 DEC 17 STN Viewer enhanced with full-text patent content  
 from USPATOLD  
 NEWS 29 JAN 02 STN pricing information for 2008 now available  
 NEWS 30 JAN 16 CAS patent coverage enhanced to include exemplified  
 prophetic substances  
 NEWS 31 JAN 28 USPATFULL, USPAT2, and USPATOLD enhanced with new  
 custom IPC display formats  
 NEWS 32 JAN 28 MARPAT searching enhanced  
 NEWS 33 JAN 28 USGENE now provides USPTO sequence data within 3 days  
 of publication  
 NEWS 34 JAN 28 TOXCENTER enhanced with reloaded MEDLINE segment  
 NEWS 35 JAN 28 MEDLINE and LEMEDLINE reloaded with enhancements  
 NEWS 36 FEB 08 STN Express, Version 8.3, now available  
  
 NEWS EXPRESS FEBRUARY 08 CURRENT WINDOWS VERSION IS V8.3,  
 AND CURRENT DISCOVER FILE IS DATED 24 JANUARY 2008  
  
 NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 17:47:21 ON 16 FEB 2008

=> File .Gerry2MBCE

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
FULL ESTIMATED COST	ENTRY	SESSION
	0.21	0.21

FILE 'MEDLINE' ENTERED AT 17:47:47 ON 16 FEB 2008

FILE 'BIOSIS' ENTERED AT 17:47:47 ON 16 FEB 2008  
Copyright (c) 2008 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 17:47:47 ON 16 FEB 2008  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 17:47:47 ON 16 FEB 2008  
Copyright (c) 2008 Elsevier B.V. All rights reserved.

=> S (MIP-4 OR CCL18 OR PARC OR AMAC1 OR AMAC-1 OR DCCK1 OR DC-CK-1 OR SCYA18 OR Ckbeta1 OR Ckbeta7) AND (CCRL2 OR HCR OR CRAM-A) AND pd<=20041202

1 FILES SEARCHED...

L1 6 (MIP-4 OR CCL18 OR PARC OR AMAC1 OR AMAC-1 OR DCCK1 OR DC-CK-1 OR SCYA18 OR CKBETA1 OR CKBETA7) AND (CCRL2 OR HCR OR CRAM-A) AND PD<=20041202

=> Dup Rem L1

PROCESSING COMPLETED FOR L1

L2 2 DUP REM L1 (4 DUPLICATES REMOVED)  
ANSWERS '1-2' FROM FILE MEDLINE

=> D Ibib abs L2 1-2

L2	ANSWER 1 OF 2	MEDLINE on STN	DUPLICATE 1
ACCESSION NUMBER:	2004617243	MEDLINE	
DOCUMENT NUMBER:	PubMed ID: 15588486		
TITLE:	Haplotype structure and linkage disequilibrium in chemokine and chemokine receptor genes.		
AUTHOR:	Clark Vanessa J; Dean Michael		
CORPORATE SOURCE:	Laboratory of Genomic Diversity, Human Genetics Section, National Cancer Institute, Frederick, MD 21702, USA.. vclark@genetics.bsd.uchicago.edu		
SOURCE:	Human genomics, (2004 May) Vol. 1, No. 4, pp. 255-73. Journal code: 101202210. ISSN: 1473-9542.		
PUB. COUNTRY:	England: United Kingdom		
DOCUMENT TYPE:	Journal; Article; (JOURNAL ARTICLE)		
LANGUAGE:	English		
FILE SEGMENT:	Priority Journals		
ENTRY MONTH:	200503		
ENTRY DATE:	Entered STN: 20 Dec 2004 Last Updated on STN: 30 Mar 2005 Entered Medline: 29 Mar 2005		

AB To dissect the haplotype structure of candidate genes for disease association studies, it is important to understand the nature of genetic variation at these loci in different populations. We present a survey of haplotype structure and linkage disequilibrium of chemokine and chemokine receptor genes in 11 geographically-distinct population samples (n=728). Chemokine proteins are involved in intercellular signalling and the immune response. These molecules are important modulators of human immunodeficiency virus (HIV)-1 infection and the progression of the acquired immune deficiency syndrome, tumour development and the metastatic process of cancer. To study the extent of genetic variation in this gene family, single nucleotide polymorphisms (SNPs) from 13 chemokine and

chemokine receptor genes were genotyped using the 5' nuclease assay (TaqMan). SNP haplotypes, estimated from unphased genotypes using the Expectation-Maximization-algorithm, are described in a cluster of four CC-chemokine receptor genes (CCR3, CCR2, CCR5 and CCR12) on chromosome 3p21, and a cluster of three CC-chemokine genes [MP1F-1 (CCL23), PARC (CCL18) and MIP-1alpha (CCL3)] on chromosome 17q11-12. The 32 base pair (bp) deletion in exon 4 of CCR5 was also included in the haplotype analysis of 3p21. A total of 87.5 per cent of the variation of 14 biallelic loci scattered over 150 kilobases of 3p21 is explained by 11 haplotypes which have a frequency of at least 1 per cent in the total sample. An analysis of haplotype blocks in this region indicates recombination between CCR2 and CCR5, although long-range pairwise linkage disequilibrium across the region appears to remain intact on two common haplotypes. A reduced-median network demonstrates a clear relationship between 3p21 haplotypes, rooted by the putative ancestral haplotype determined by direct sequencing of four primate species. Analysis of six SNPs on 17q11-12 indicates that 97.5 per cent of the variation is explained by 15 haplotypes, representing at least 1 per cent of the total sample. Additionally, a possible signature of selection at a non-synonymous coding SNP (M106V) in the MP1F-1 (CCL23) gene warrants further study. We anticipate that the results of this study of chemokine and chemokine receptor variation will be applicable to more extensive surveys of long-range haplotype structure in these gene regions and to association studies of HIV-1 disease and cancer.

L2	ANSWER 2 OF 2	MEDLINE on STN	DUPLICATE 2
ACCESSION NUMBER:	2004617237	MEDLINE	
DOCUMENT NUMBER:	PubMed ID: 15588479		
TITLE:	Characterisation of SNP haplotype structure in chemokine and chemokine receptor genes using CEPH pedigrees and statistical estimation.		
AUTHOR:	Clark Vanessa J; Dean Michael		
CORPORATE SOURCE:	Laboratory of Genomic Diversity, Human Genetics Section, National Cancer Institute, Frederick, MD 21702, USA.. vclark@genetics.bsd.uchicago.edu		
SOURCE:	Human genomics, (2004 Mar) Vol. 1, No. 3, pp. 195-207. Journal code: 101202210. ISSN: 1473-9542.		
PUB. COUNTRY:	England; United Kingdom		
DOCUMENT TYPE:	Journal; Article; (JOURNAL ARTICLE)		
LANGUAGE:	English		
FILE SEGMENT:	Priority Journals		
ENTRY MONTH:	200501		
ENTRY DATE:	Entered STN: 20 Dec 2004 Last Updated on STN: 19 Jan 2005 Entered Medline: 18 Jan 2005		
AB	Chemokine signals and their cell-surface receptors are important modulators of HIV-1 disease and cancer. To aid future case/control association studies, aim to further characterise the haplotype structure of variation in chemokine and chemokine receptor genes. To perform haplotype analysis in a population-based association study, haplotypes must be determined by estimation, in the absence of family information or laboratory methods to establish phase. Here, test the accuracy of estimates of haplotype frequency and linkage disequilibrium by comparing estimated haplotypes generated with the expectation maximisation (EM) algorithm to haplotypes determined from Centre d'Etude Polymorphisme Humain (CEPH) pedigree data. To do this, they have characterised haplotypes comprising alleles at 11 biallelic loci in four chemokine receptor genes (CCR3, CCR2, CCR5 and CCR12), which span 150 kb on chromosome 3p21, and haplotypes of nine biallelic loci in six chemokine genes [MCP-1(CCL2), Eotaxin(CCL11), RANTES(CCL5), MP1F-1(CCL23), PARC(CCL18) and MIP-1alpha(CCL3)] on chromosome		

17q11-12. Forty multi-generation CEPH families, totalling 489 individuals, were genotyped by the TaqMan 5'-nuclease assay. Phased haplotypes and haplotypes estimated from unphased genotypes were compared in 103 grandparents who were assumed to have mated at random. For the 3p21 single nucleotide polymorphism (SNP) data, haplotypes determined by pedigree analysis and haplotypes generated by the EM algorithm were nearly identical. Linkage disequilibrium, measured by the D' statistic, was nearly maximal across the 150 kb region, with complete disequilibrium maintained at the extremes between CCR3-Y17Y and CCRL2-I243V. D'-values calculated from estimated haplotypes on 3p21 had high concordance with pairwise comparisons between pedigree-phased chromosomes. Conversely, there was less agreement between analyses of haplotype frequencies and linkage disequilibrium using estimated haplotypes when compared with pedigree-phased haplotypes of SNPs on chromosome 17q11-12. These results suggest that, while estimations of haplotype frequency and linkage disequilibrium may be relatively simple in the 3p21 chemokine receptor cluster in population samples, the more complex environment on chromosome 17q11-12 will require a higher resolution haplotype analysis.

```
=> log off H
SESSION WILL BE HELD FOR 120 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 17:49:55 ON 16 FEB 2008
```

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAEGS1646

PASSWORD:  
 TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

```
NEWS 1      Web Page for STN Seminar Schedule - N. America
NEWS 2 AUG 06 CAS REGISTRY enhanced with new experimental property tags
NEWS 3 AUG 06 FSTA enhanced with new thesaurus edition
NEWS 4 AUG 13 CA/CAPlus enhanced with additional kind codes for granted
           patents
NEWS 5 AUG 20 CA/CAPlus enhanced with CAS indexing in pre-1907 records
NEWS 6 AUG 27 Full-text patent databases enhanced with predefined
           patent family display formats from INPADOCDB
NEWS 7 AUG 27 USPATOLD now available on STN
NEWS 8 AUG 28 CAS REGISTRY enhanced with additional experimental
           spectral property data
NEWS 9 SEP 07 STN AnaVist, Version 2.0, now available with Derwent
           World Patents Index
NEWS 10 SEP 13 FORIS renamed to SOFIS
NEWS 11 SEP 13 INPADOCDB enhanced with monthly SDI frequency
NEWS 12 SEP 17 CA/CAPlus enhanced with printed CA page images from
           1967-1998
NEWS 13 SEP 17 CAPlus coverage extended to include traditional medicine
           patents
NEWS 14 SEP 24 EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS 15 OCT 02 CA/CAPlus enhanced with pre-1907 records from Chemisches
           Zentralblatt
NEWS 16 OCT 19 BEILSTEIN updated with new compounds
```



NEWS 17 NOV 15 Derwent Indian patent publication number format enhanced  
 NEWS 18 NOV 19 WPIX enhanced with XML display format  
 NEWS 19 NOV 30 ICSD reloaded with enhancements  
 NEWS 20 DEC 04 LINPADOCDB now available on STN  
 NEWS 21 DEC 14 BEILSTEIN pricing structure to change  
 NEWS 22 DEC 17 USPATOLD added to additional database clusters  
 NEWS 23 DEC 17 IMSDRUGCONF removed from database clusters and STN  
 NEWS 24 DEC 17 DGENE now includes more than 10 million sequences  
 NEWS 25 DEC 17 TOXCENTER enhanced with 2008 MeSH vocabulary in  
 MEDLINE segment  
 NEWS 26 DEC 17 MEDLINE and LMEDLINE updated with 2008 MeSH vocabulary  
 NEWS 27 DEC 17 CA/Caplus enhanced with new custom IPC display formats  
 NEWS 28 DEC 17 STN Viewer enhanced with full-text patent content  
 from USPATOLD  
 NEWS 29 JAN 02 STN pricing information for 2008 now available  
 NEWS 30 JAN 16 CAS patent coverage enhanced to include exemplified  
 prophetic substances  
 NEWS 31 JAN 28 USPATFULL, USPAT2, and USPATOLD enhanced with new  
 custom IPC display formats  
 NEWS 32 JAN 28 MARPAT searching enhanced  
 NEWS 33 JAN 28 USGENE now provides USPTO sequence data within 3 days  
 of publication  
 NEWS 34 JAN 28 TOXCENTER enhanced with reloaded MEDLINE segment  
 NEWS 35 JAN 28 MEDLINE and LMEDLINE reloaded with enhancements  
 NEWS 36 FEB 08 STN Express, Version 8.3, now available

NEWS EXPRESS FEBRUARY 08 CURRENT WINDOWS VERSION IS V8.3,  
 AND CURRENT DISCOVER FILE IS DATED 24 JANUARY 2008

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
 NEWS LOGIN Welcome Banner and News Items  
 NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that  
 specific topic.

All use of STN is subject to the provisions of the STN Customer  
 agreement. Please note that this agreement limits use to scientific  
 research. Use for software development or design or implementation  
 of commercial gateways or other similar uses is prohibited and may  
 result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 20:11:55 ON 16 FEB 2008

=> File .Gerry2MBCE	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 20:12:20 ON 16 FEB 2008

FILE 'BIOSIS' ENTERED AT 20:12:20 ON 16 FEB 2008  
 Copyright (c) 2008 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 20:12:20 ON 16 FEB 2008  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> S MIP-4 (S)CCRL2  
 L1 1 MIP-4 (S) CCRL2

=> D abs

L1 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN  
 AB Macrophage inflammatory protein-4 (MIP-4; also known as DC-CK1, CCL18 and PARC) is identified as an endogenous ligand for CCRL2 (chemokine (C-C motif) receptor-like 2). The protein and cDNA sequences of human MIP-4 and CCRL2 are disclosed. Anti-CCRL2 antibody was blocking MIP-4 and synovial fluid induced monocyte chemotaxis. Anti-MIP-4 antibody was also blocking RA synovial fluid induced monocyte chemotaxis. This data demonstrates that MIP-4 is a major mediator of monocyte induced chemotaxis found in RA synovial fluid. CCRL2 modulators, such as antibodies against CCRL2 or MIP-4, is useful in treating an inflammatory disease, a disease associated with enhanced macrophage activity or an infection.

=> D Ibib

L1 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN  
 ACCESSION NUMBER: 2005:547792 CAPLUS  
 DOCUMENT NUMBER: 143:76842  
 TITLE: Macrophage inflammatory protein-4 (MIP-4) as an endogenous ligand for CCRL2 and sequences of human MIP-4 and CCRL2  
 INVENTOR(S): Tinsley, Jonathon Mark  
 PATENT ASSIGNEE(S): Oxagen Limited, UK  
 SOURCE: PCT Int. Appl., 69 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005057220	A2	20050623	WO 2004-GB5057	20041202
WO 2005057220	A3	20060202		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1692171	A2	20060823	EP 2004-801256	20041202
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, BA, HR, IS, YU				
JP 2007520210	T	20070726	JP 2006-542005	20041202
US 2007036781	A1	20070215	US 2006-579386	20060515

PRIORITY APPLN. INFO.:

GB 2003-28275	A	20031205
GB 2004-3014	A	20040211
GB 2004-18568	A	20040819
WO 2004-GB5057	W	20041202

=> S MIP-4 (S) receptor  
L2 4 MIP-4 (S) RECEPTOR

=> Dup Rem L2  
PROCESSING COMPLETED FOR L2  
L3 4 DUP REM L2 (0 DUPLICATES REMOVED)  
ANSWER '1' FROM FILE BIOSIS  
ANSWERS '2-4' FROM FILE CAPLUS

=> D Ibib abs L3 1-4

L3 ANSWER 1 OF 4 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN  
ACCESSION NUMBER: 2007:307109 BIOSIS  
DOCUMENT NUMBER: PREV200700295995  
TITLE: Histamine release from the basophils of control and  
asthmatic subjects and a comparison of gene expression  
between "releaser". and "nonreleaser" basophils.  
AUTHOR(S): Youssef, Lama A.; Schuyler, Mark; Gilmartin, Laura;  
Pickett, Gavin; Bard, Julie D. J.; Tarleton, Christy A.;  
Archibeque, Tereassa; Qualls, Clifford; Wilson, Bridget S.;  
Oliver, Janet M. [Reprint Author]  
CORPORATE SOURCE: Univ New Mexico, Sch Med, Dept Cell Pathol Lab, 2325 Camino  
de Salud, Albuquerque, NM 87131 USA  
joliver@salud.unm.edu  
SOURCE: Journal of Immunology, (APR 1 2007) Vol. 178, No. 7, pp.  
4584-4594.  
CODEN: JOIMA3. ISSN: 0022-1767.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 9 May 2007  
Last Updated on STN: 9 May 2007

AB Most human blood basophils respond to Fc epsilon RI cross-linking by releasing histamine and other inflammatory mediators. Basophils that do not degranulate after anti-IgE challenge, known as "nonreleaser" basophils, characteristically have no or barely detectable levels of the Syk tyrosine kinase. The true incidence of the nonreleaser phenotype, its relationship (if any) to allergic asthma, and its molecular mechanism are not well understood. In this study, we report statistical analyses of degranulation assays performed in 68 control and 61 asthmatic subjects that establish higher basal and anti-IgE-stimulated basophil degranulation among the asthmatics. Remarkably, 28% of the, control group and 13% of the asthmatic group were nonreleasers; for all or part of our 4-year long study and cycling between the releaser and nonreleaser phenotypes occurred at least once in blood basophils from 8 (of 8) asthmatic and 16 (of 23) control donors. Microarray analysis showed that basal gene expression was generally lower in nonreleaser than releaser basophils. In releaser I cells, Fc epsilon RI cross-linking up-regulated > 200 genes, including genes encoding receptors (the Fc epsilon RI alpha and beta subunits, the histamine 4 receptor, the chemokine (C-C motif) receptor 1), signaling proteins (Lyn), chemokines (IL-8, RANTES, MIP-1 alpha, and, MIP-4 beta) and transcription factors (early growth response-1, early growth response-3, and AP-1). Fc epsilon RI cross-linking induced fewer, and quite distinct, transcriptional responses in nonreleaser cells. We conclude that "nonreleaser" and "cycler" basophils represent a distinct and reversible natural phenotype. Although histamine is more readily released from

basophils isolated from asthmatics than controls, the presence of  
noureleaser basophils does not rule out the diagnosis of asthma.

L3 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:547792 CAPLUS

DOCUMENT NUMBER: 143:76842

TITLE: Macrophage inflammatory protein-4 (MIP-4) as an  
endogenous ligand for CCRL2 and sequences of human  
MIP-4 and CCRL2

INVENTOR(S): Tinsley, Jonathon Mark

PATENT ASSIGNEE(S): Oxagen Limited, UK

SOURCE: PCT Int. Appl., 69 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005057220	A2	20050623	WO 2004-GB5057	20041202
WO 2005057220	A3	20060202		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1692171	A2	20060823	EP 2004-801256	20041202
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, BA, HR, IS, YU			
JP 2007520210	T	20070726	JP 2006-542005	20041202
US 2007036781	A1	20070215	US 2006-579386	20060515
PRIORITY APPLN. INFO.:			GB 2003-28275	A 20031205
			GB 2004-3014	A 20040211
			GB 2004-18568	A 20040819
			WO 2004-GB5057	W 20041202
AB	Macrophage inflammatory protein-4 (MIP-4; also known as DC-CKI, CCL18 and PARC) is identified as an endogenous ligand for CCRL2 (chemokine (C-C motif) receptor-like 2). The protein and cDNA sequences of human MIP-4 and CCRL2 are disclosed. Anti-CCRL2 antibody was blocking MIP-4 and synovial fluid induced monocyte chemotaxis. Anti-MIP-4 antibody was also blocking RA synovial fluid induced monocyte chemotaxis. This data demonstrates that MIP-4 is a major mediator of monocyte induced chemotaxis found in RA synovial fluid. CCRL2 modulators, such as antibodies against CCRL2 or MIP-4, is useful in treating an inflammatory disease, a disease associated with enhanced macrophage activity or an infection.			
L3 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN				
ACCESSION NUMBER:	2000:82601 CAPLUS			
DOCUMENT NUMBER:	132:221179			
TITLE:	C-C chemokine receptor 3 antagonism by the β-chemokine macrophage inflammatory protein 4, a property strongly enhanced by an amino-terminal alanine-methionine swap			

AUTHOR(S): Nibbs, Robert J. B.; Salcedo, Theodora W.; Campbell, John D. M.; Yao, Xiao-Tao; Li, Yuling; Nardelli, Bernardetta; Olsen, Henrik S.; Morris, Tina S.; Proudfoot, Amanda E. I.; Patel, Vikram P.; Graham, Gerard J.

CORPORATE SOURCE: Cancer Research Campaign Laboratories, Beatson Institute for Cancer Research, Glasgow, G61 1BD, UK

SOURCE: Journal of Immunology (2000), 164(3), 1488-1497  
CODEN: JOIMA3; ISSN: 0022-1767

PUBLISHER: American Association of Immunologists

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Allergic reactions are characterized by the infiltration of tissues by activated eosinophils, Th2 lymphocytes, and basophils. The  $\beta$ -chemokine receptor CCR3, which recognizes the ligands eotaxin, eotaxin-2, monocyte chemoattractant protein (MCP) 3, MCP4, and RANTES, plays a central role in this process, and antagonists to this receptor could have potential therapeutic use in the treatment of allergy. The authors describe here a potent and specific CCR3 antagonist, called Met-chemokine  $\beta$  7 (Ck $\beta$ 7), that prevents signaling through this receptor and, at concns. as low as 1 nM, can block eosinophil chemotaxis induced by the most potent CCR3 ligands. Met-Ck $\beta$ 7 is a more potent CCR3 antagonist than Met- and aminooxypentane (AOP)-RANTES and, unlike these proteins, exhibits no partial agonist activity and is highly specific for CCR3. This antagonist may thus be of use in ameliorating leukocyte infiltration associated with allergic inflammation. Met-Ck $\beta$ 7 is a modified form of the  $\beta$ -chemokine macrophage inflammatory protein (MIP) 4 [alternatively called pulmonary and activation-regulated chemokine (PARC), alternative macrophage activation-associated C-C chemokine (AMAC) 1, or dendritic cell-derived C-C chemokine (DCCK) 1]. Surprisingly, the unmodified MIP4 protein, which is known to act as a T cell chemoattractant, also exhibits this CCR3 antagonistic activity, although to a lesser extent than Met-Ck $\beta$ 7, but to a level that may be of physiolo. relevance. MIP4 may therefore use chemokine receptor agonism and antagonism to control leukocyte movement in vivo. The enhanced activity of Met-Ck $\beta$ 7 is due to the alteration of the extreme N-terminal residue from an alanine to a methionine.

REFERENCE COUNT: 53 THERE ARE 53 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1995:820779 CAPLUS

DOCUMENT NUMBER: 123:220290

TITLE: Cloning and therapeutic applications of human macrophage inflammatory proteins MIP-3, MIP-4, and MIP-1y, or their antibodies or antagonists

INVENTOR(S): Li, Haodong; Rosen, Craig A.; Ruben, Steve; Adams, Mark D.

PATENT ASSIGNEE(S): Human Genome Sciences, Inc., USA

SOURCE: PCT Int. Appl., 63 pp.  
CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 10

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9517092	A1	19950629	WO 1994-US7256	19940628
W: AU, CA, CN, JP, KR, NZ				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5556767	A	19960917	US 1993-173209	19931222

US 5504003	A	19960402	US 1994-208339	19940308
ZA 9403442	A	19951120	ZA 1994-3442	19940518
CA 2179606	A1	19950629	CA 1994-2179606	19940628
AU 9475497	A	19950710	AU 1994-75497	19940628
AU 684539	B2	19971218		
EP 735818	A1	19961009	EP 1994-925671	19940628
EP 735818	B1	20040331		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
CN 1143894	A	19970226	CN 1994-194902	19940628
JP 09506774	T	19970708	JP 1995-517397	19940628
JP 3677288	B2	20050727		
JP 2002053490	A	20020219	JP 2001-196723	19940628
AT 262914	T	20040415	AT 1994-925671	19940628
PT 735818	T	20040730	PT 1994-925671	19940628
ES 2214484	T3	20040916	ES 1994-925671	19940628
CN 1321745	A	20011114	CN 2001-116577	20010416
AU 777297	B2	20041007	AU 2002-15445	20020206
US 2003147846	A1	20030807	US 2002-165233	20020610

PRIORITY APPLN. INFO.:

US 1993-173209	A	19931222
US 1994-208339	A	19940308
JP 1995-517397	A3	19940628
WO 1994-US7256	W	19940628
US 1995-446881	B1	19950505
US 1995-468775	B2	19950606
AU 1997-46576	A3	19970930
US 1999-334923	A3	19990617
US 1999-334951	A3	19990617
US 1999-334954	A3	19990617

AB There are disclosed human macrophage inflammatory protein-3, human macrophage inflammatory protein-4, and human macrophage inflammatory protein-ly polypeptides and DNA (or RNA) encoding such polypeptides. There is also provided a procedure for producing such polypeptides by recombinant techniques and for producing antibodies against such polypeptides. In the invention there is also provided antagonist/inhibitors against such polypeptides which inhibit the functioning of such polypeptides. Another aspect of the invention provides a combination of the polypeptides of the present invention and a suitable pharmaceutical carrier for providing a therapeutically effective amount of the polypeptides for the treatment of various associated diseases.

=> S CCRL2 (S) ligand  
L4 5 CCRL2 (S) LIGAND

=> Dup Rem L4  
PROCESSING COMPLETED FOR L4  
L5 3 DUP REM L4 (2 DUPLICATES REMOVED)  
ANSWER '1' FROM FILE MEDLINE  
ANSWER '2' FROM FILE BIOSIS  
ANSWER '3' FROM FILE CAPLUS

=> D Ibib Abs L5 1-3

L5 ANSWER 1 OF 3	MEDLINE on STN	DUPLICATE 1
ACCESSION NUMBER:	2004286185	MEDLINE
DOCUMENT NUMBER:	PubMed ID: 15188357	
TITLE: Up-regulated expression and activation of the orphan chemokine receptor, CCRL2, in rheumatoid arthritis.		
AUTHOR: Galligan Carole L; Matsuyama Wataru; Matsukawa Akihiro; Mizuta Hiroshi; Hodge David R; Howard O M Zack; Yoshimura Teizo		
CORPORATE SOURCE: National Cancer Institute at Frederick, Frederick, Maryland		

21702, USA.  
 SOURCE: Arthritis and rheumatism, (2004 Jun) Vol. 50, No. 6, pp. 1806-14.  
 Journal code: 0370605. ISSN: 0004-3591.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals  
 ENTRY MONTH: 200407  
 ENTRY DATE: Entered STN: 10 Jun 2004  
 Last Updated on STN: 9 Jul 2004  
 Entered Medline: 8 Jul 2004

AB OBJECTIVE: Rheumatoid arthritis (RA) is a chronic inflammatory condition characterized by a cellular influx and destruction of the joint architecture. Chemokines characteristically regulate leukocyte recruitment and activation. Chemokine (CC motif) receptor-like 2 (CCRL2) is an orphan receptor with homology to other CC chemokine receptors. We undertook this study to examine CCRL2 expression in RA, cytokine regulation of expression, and the source of a putative ligand in an attempt to determine the role of this receptor during inflammation. METHODS: Expression of CCRL2 on joint-infiltrating leukocytes was examined by immunocytochemistry. In vitro studies evaluated CCRL2 expression in primary neutrophils using Northern and Western blotting and reverse transcriptase-polymerase chain reaction. HEK 293 cells expressing two splice variants of CCRL2 (HEK/CCRL2A or HEK/CCRL2B) were generated with a retroviral expression system, and their migration in response to fractions of synovial fluid (SF) from RA patients was examined using a 48-well chamber. RESULTS: CCRL2 expression was observed on all infiltrating neutrophils and on some macrophages obtained from the SF of 5 RA patients. In vitro studies of primary neutrophils revealed that CCRL2 messenger RNA (mRNA) was rapidly up-regulated following stimulation with lipopolysaccharide (1 microg/ml) or tumor necrosis factor (5 ng/ml). The mRNA for both CCRL2A and CCRL2B were expressed in cytokine-stimulated neutrophils. Cells expressing either of these splice variants migrated in response to a fraction of RA SF. CONCLUSION: CCRL2 expression is up-regulated on synovial neutrophils of RA patients. Inflammatory products present in the SF activate this receptor, indicating that CCRL2 is a functional receptor that may be involved in the pathogenesis of RA.

L5 ANSWER 2 OF 3 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN  
 ACCESSION NUMBER: 2004:286307 BIOSIS  
 DOCUMENT NUMBER: PREV200400285064  
 TITLE: Upregulated expression and activation of the orphan chemokine receptor, CCRL2, in rheumatoid arthritis.  
 AUTHOR(S): Galligan, Carole [Reprint Author]; Matsuyama, Wataru; Matsukawa, Akihiro; Mizuta, Hiroshi; Hodge, David R; Howard, O.M. Zack; Yoshimura, Teizo  
 CORPORATE SOURCE: Laboratory of Molecular Immunoregulation, National Cancer Institute, P.O. Box B, Bldg. 560, Frederick, MD, 21702-1201, USA  
 cgalligan@ncicfcrf.gov  
 SOURCE: FASEB Journal, (2004) Vol. 18, No. 4-5, pp. Abst. 337.9.  
<http://www.fasebj.org/>. e-file.  
 Meeting Info.: FASEB Meeting on Experimental Biology: Translating the Genome. Washington, District of Columbia, USA. April 17-21, 2004. FASEB.  
 ISSN: 0892-6638 (ISSN print).  
 DOCUMENT TYPE: Conference; (Meeting)  
 Conference; Abstract; (Meeting Abstract)  
 LANGUAGE: English  
 ENTRY DATE: Entered STN: 16 Jun 2004  
 Last Updated on STN: 16 Jun 2004

AB Rheumatoid arthritis (RA) is a chronic inflammatory condition characterized by a cellular influx and destruction of the joint architecture involving chemokines that induce the leukocyte infiltration and activation. The human chemokine-like receptor 2 (CCRL2) codes for a putative 7-TM G protein-coupled receptor with high homology to other chemokine receptors. This study examined CCRL2 expression in RA, the cytokines regulating gene expression and the source of a putative ligand for CCRL2 in an attempt to determine the role of this receptor during inflammation. Immunohistochemistry revealed positive CCRL2 staining of neutrophils infiltrating the joints of RA patients. Primary human neutrophils expressed low levels of CCRL2 mRNA, but stimulation with LPS or TNF dramatically upregulated mRNA levels. Elevated CCRL2 mRNA expression was evident as early as 1 h after TNF- or LPS-activation and the levels peaked after 2-4 or 4-8 hours respectively. Two N-terminal splice variants for CCRL2 (A and B) were detected in freshly isolated as well as in LPS- and TNF-activated neutrophils by RT-PCR. CCRL2 protein was not detectable in freshly isolated neutrophils but readily detectable in LPS-activated neutrophils. Fractions of RA synovial fluids induced significant chemotaxis for HEK-293 cells expressing either CCRL2 variant. Our results suggest that CCRL2 may play a role in regulating neutrophil recruitment and activation during rheumatoid arthritis.

L5 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2008 ACS on SIN

ACCESSION NUMBER: 2005:547792 CAPLUS  
DOCUMENT NUMBER: 143:76842  
TITLE: Macrophage inflammatory protein-4 (MIP-4) as an endogenous ligand for CCRL2 and sequences of human MIP-4 and CCRL2  
INVENTOR(S): Tinsley, Jonathon Mark  
PATENT ASSIGNEE(S): Oxagen Limited, UK  
SOURCE: PCT Int. Appl., 69 pp.  
CODEN: P1XXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005057220	A2	20050623	WO 2004-GB5057	20041202
WO 2005057220	A3	20060202		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1692171	A2	20060823	EP 2004-801256	20041202
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, BA, HR, IS, YU			
JP 2007520210	T	20070726	JP 2006-542005	20041202
US 2007036781	A1	20070215	US 2006-579386	20060515
PRIORITY APPLN. INFO.:			GB 2003-28275	A 20031205
			GB 2004-3014	A 20040211
			GB 2004-18568	A 20040819



AB Macrophage inflammatory protein-4 (MIP-4; also known as DC-CKI, CCL18 and PARC) is identified as an endogenous ligand for CCRL2 (chemokine (C-C motif) receptor-like 2). The protein and cDNA sequences of human MIP-4 and CCRL2 are disclosed. Anti-CCRL2 antibody was blocking MIP-4 and synovial fluid induced monocyte chemotaxis. Anti-MIP-4 antibody was also blocking RA synovial fluid induced monocyte chemotaxis. This data demonstrates that MIP-4 is a major mediator of monocyte induced chemotaxis found in RA synovial fluid. CCRL2 modulators, such as antibodies against CCRL2 or MIP-4, is useful in treating an inflammatory disease, a disease associated with enhanced macrophage activity or an infection.

=> Log Off H

SESSION WILL BE HELD FOR 120 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 20:19:13 ON 16 FEB 2008